

Bluetongue in sheep Know the signs

Bluetongue serotype 3 (BTV-3) is circulating on the eastern side of England. The Animal and Plant Health Agency (APHA) and Defra have declared bluetongue zones spanning several counties to help control further spread of the disease. You can find out more and check whether you are in a bluetongue zone by visiting gov.uk/bluetongue.

You can play a part in helping to stop bluetongue spreading by checking your animals for clinical signs, only moving animals if essential.



- · ulcers or sores in the mouth and nose
- discharge from the eyes or nose and drooling from the mouth
- swelling of the lips, tongue, head and neck
- heat and tenderness at the coronary band (where the skin of the leg meets the horn of the foot)

Other clinical signs include:

- inappetence and weight loss
- red skin as a result of blood collecting beneath the surface
- fever
- · lameness and reluctance to move
- breathing problems
- abortion, foetal deformities and stillbirths
- death

In lambs

Lambs can become infected with bluetongue virus before birth if the dam is infected while pregnant. Signs of infection include:

- · lambs born small, weak, deformed or blind
- · death of lambs within a few days of birth
- stillbirths



Ulcers and lesions in the mouth



Nasal redness



Nasal discharge



Recumbant animal

Find out more information and how to report bluetongue by visiting www.gov.uk/bluetongue

